Pre-Calculus 11 HW 4.4 The Quadratic Formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad , \quad a \neq 0$$

1. Given each quadratic equation, indicate the values of the coefficients "a", "b" and "c":

a) $x^2 + 5x + 6 = 0$	b) $12x^2 + 7x - 3 = 0$	c) $-2x^2 - 7x + 5 = 0$
<i>a</i> = <i>b</i> = <i>c</i> =	a = b = c =	<i>a</i> = <i>c</i> =
d) $4x^2 = 13x - 8$	e) $x(7-8x) = 10$	f) $x(x+2) = 6 - (x-3)(2x+1)$
a = b = c =	a =b =c =	a =b =c =

2. Solve for "x" for each of the following by using the quadratic formula:

a) $x^2 - 5x + 6 = 0$	b) $3x^2 - x + 1 = 3$	c) $2x^2 - 3x - 1 = 0$
d) $-0.5x^2 + 4x + 12 = 0$	e) $-0.5x^2 + 3x = 6$	f) $-3x^2 = 12x - 5$

h) $\frac{4x^2}{3} = 4x - 2$	i) $x^2 - 2 = \frac{-7x}{2}$
k) $2x^2 + 6x - 8 = 7x^2 - 2x$	L) $2(2x-1)^2 + 9(2x-1) + 7 = 0$

- 3. Under what conditions will the equation have no solutions? $ax^2 + bx + c = 0$ Explain why
- 4. Under what conditions will the equation have only one solution? $ax^2 + bx + c = 0$ Explain why
- 5. The revenue a company makes for selling shoes is given by the equation: $R = -8p^2 + 1200p$, where "R" is the revenue and 'p" is the price in dollars. At what price should the company sell their shoes to generate a revenue of \$400,000?

6. Here are the steps that John used to solve the equation: $12x^2 - 7x - 3 = 0$. Find the mistakes:

$$L_1: x = -7 \pm \frac{\sqrt{49^2 - 4(12)(3)}}{2(12)}$$

$$L_4: x = -7 \pm 1.9794955..$$

$$L_2: x = -7 \pm \frac{\sqrt{2401 - 144}}{24}$$

$$L_5: x_1 = -7 + 1.9794955.. = -5.0205...$$

$$L_3: x = -7 \pm \frac{\sqrt{2257}}{24}$$

$$L_6: x_2 = -7 - 1.9794955... = -8.9794955...$$