Data	
Date	

Pre-Calculus 11 CH3/4 HW Lesson 5: The Quadratic Formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad , \ a \neq 0$$

1. Given each quadratic equation, indicate the values of the coefficients "a", "b" and "c":

a)
$$x^2 + 5x + 6 = 0$$
 b) $12x^2 + 7x - 3 = 0$
 c) $-2x^2 - 7x + 5 = 0$
 $a = __b = __c = __a$
 $a = __b = __c = __a$
 $a = __b = __c = __a$
 $d) 4x^2 = 13x - 8$
 e) $x(7 - 8x) = 10$
 f) $x(x + 2) = 6 - (x - 3)(2x + 1)$
 $a = __b = __c = __a$
 $a = __b = __c = __a$
 $a = __b = __c = __a$

2. Solve for "x" for each of the following by using the quadratic formula:

a) $x^2 - 5x + 6 = 0$	b) $3x^2 - x + 1 = 3$	c) $2x^2 - 3x - 1 = 0$
d) $-0.5x^2 + 4x + 12 = 0$	e) $-0.5x^2 + 3x = 6$	f) $-3x^2 = 12x - 5$

g) $3x(2x-6) = 8$	h) $\frac{4x^2}{3} = 4x - 2$	i) $x^2 - 2 = \frac{-7x}{2}$
j) $2x^3 - 5x^2 + 7x = 0$	k) $2x^2 + 6x - 8 = 7x^2 - 2x$	L) $2(2x-1)^2 + 9(2x-1) + 7 = 0$

3. Under what conditions will the equation have no solutions? $ax^2 + bx + c = 0$ Explain why

- 4. Under what conditions will the equation have only one solution? $ax^2 + bx + c = 0$ Explain why
- 5. The revenue a company makes for selling shoes is given by the equation: $R = -8p^2 + 1200p$, where "*R*" is the revenue and 'p" is the price in dollars. At what price should the company sell their shoes to generate a revenue of \$400,000?

6. Here are the steps that John used to solve the equation: $12x^2 - 7x - 3 = 0$. Find the mistakes:

Copyright All Rights Reserved at www.BCMath.ca